

Selected flavonoids and their role in the treatment of epilepsy – a review of the latest reports from experimental studies

Joanna Szala-Rycaj*, Mirosław Zagaja, Aleksandra Szewczyk and Marta Andres-Mach

Isobolographic Analysis Laboratory, Institute of Rural Health, Lublin, Poland,

** Email: joanna.szala@onet.pl*

Epilepsy is a chronic neurological disease characterized by recurrent seizures that affects about 70 million people worldwide. Antiepileptic drugs are the most commonly used medications in the treatment of epilepsy. They help control seizures in about 60-70% of people. The remaining percentage of patients suffer from drug-resistant epilepsy, prompting scientists to look for natural substances that would prevent seizures or support the effects of drugs in add-on therapy while reducing side effects. Currently, there is a lot of emphasis on natural product. Flavonoids are included in this group, and their use in the treatment of epilepsy could support the effect of other drugs. Due to very good results of preclinical studies, flavonoids are a promising candidate for epilepsy related clinical trials related. The article is an overview of literature reports from the past 10 years including mainly *in vivo* preclinical research on various models of experimental epilepsy with the use of selected flavonoids.

Key words: flavonoids, antiepileptic drugs, experimental epilepsy

INTRODUCTION

Epilepsy is a serious neurological disorder affecting around 70 million people around the world. Antiepileptic drug (AED) therapy is effective in 60–70% of patients, however in 30-40% of patients seizure control is not achieved (Kalilani et al., 2018). Drug resistance in epilepsy is defined by International League Against Epilepsy as the failure of treatment with two properly used and well-tolerated AEDs (both, monotherapy and combination) used for a sufficiently long time in controlling seizures (Kwan et al., 2010). The mechanism of drug resistance is not fully known or fully explained. It is probably influenced by both genetic and environmental factors (Tang et al., 2017). For those suffering from drug-resistant epilepsy (DRE) numerous drug side effects, memory disorders, stigmatization become a social problem, and the lack of expected treatment effects contribute to the occurrence of depression in

even half of the patients (Kozera-Kępnia et al., 2013). Sudden unexpected death in epilepsy (SUDEP) is the main cause of death in patients with DRE and accounts for up to 50% documented causes of death in this group of patients (Opeskin and Berkovic 2003; Tomson et al., 2008; Surges et al., 2009; Shorvon and Tomson, 2011). Therefore, the main goal of DRE therapy is to extend the seizure control period, minimize side effects and improve quality of life (Boon et al., 2015). Currently, a number of scientific studies include the use of natural substances with antiepileptic properties as adjuncts to standard antiepileptic drugs in the DRE treatment and flavonoids and may be a potential group of natural drugs that increase the effects of commonly used anti-convulsants.

Flavonoids, a class of secondary plant metabolites, have been shown to demonstrate a wide spectrum of biological activity, among others antioxidant (Santos et al., 1999), anti-inflammatory (Pintho et al., 2016), anti-

allergic (Kawai et al., 2006), antiatherosclerotic (Goldwasser et al., 2011), antidiabetic (Rayidi et al., 2011) as well as hepatoprotective (Renugadevi et al., 2009). Corsale et al. (2018) used a mixture of flavonoids such as diosmin, troxerutin, rutin, hesperidin, and quercetin to reduce bleeding due to hemorrhoidal disease. Cicero et al. (2019) evaluated the effectiveness of bergamot flavonoid extract in people with dyslipidemia and those overweight. Obtained results showed an improvement in lipid and glucose metabolism, adipokine pattern and systemic inflammation in overweight and dyslipidemic individuals. Kirienko et al. (2018) investigated the clinical efficacy of 1000 mg micronized purified flavonoid fraction (MPFF) administered once daily in patients with chronic venous disease (CVD), which turned out to be associated with a rapid reduction in leg pain throughout the 8-week treatment period.

Flavonoids are derivatives of 2-phenyl-benzo- γ -pyrone. A common part in the chemical structure of all flavonoids is the carbon skeleton based on the flavane system (C6-C3-C6), formed from two benzene rings (A and B) connected by a heterocyclic ring of pyran or pyrone (C) (Fig. 1, Pietta, 2000). The biosynthesis of A and B rings occurs in two pathways – acetate (A) and shikimate (B). Ring A is formed from 3 malonyl-CoA molecules obtained from glucose conversions. Ring B is formed from 4-coumaroyl-CoA, created in the shikimate pathway from phenylalanine. Condensation of the A and B ring leads to the formation of chalcone, which is cyclized with the participation of isomerase and forms flavanone – the initial compound for the synthesis of the remaining flavonoid groups (Bravo et al., 1998, Wang et al., 2018). They may exist in free form of aglycone or betaglycosides (Panche et al., 2016). Due to differences in the structure, flavonoid compounds are divided into several subclasses: flavanones, flavonols, flavones, isoflavones, chalcones and anthocyanins (Table I.) The article focuses on several compounds from the flavonoid group that have been shown to possess




Fig. 1. The main structure of flavonoids with rings and numbered positions.

antiepileptic properties in various experimental models of epilepsy (Table II).

Anticonvulsant properties of flavonoids

Naringenin/naringin

Naringenin (2,3-dihydro-5,7-dihydroxy-2-(4-hydroxyphenyl)-4H-1-benzopyran-4-one, Fig. 2) is a natural phenolic compound found in citrus species and Chinese plants such as: *Drynaria fortunei*, *Citrus aurantium* and *Citrus medica* (Jadeja et al., 2014; Chen et al., 2016). Shakeel et al. (2017) indicated anticonvulsant properties for naringenin in the mouse pilocarpine-induced seizures (PILO). Results they obtained showed that 15-day treatment naringine (20 mg/kg and 40 mg/kg), followed by the administration of pilocarpine on the last day of treatment, seizures reduced in comparison with the control group. Naringenin has been shown to inhibit the onset and duration of seizures in a dose-dependent manner. Naringin in a dose of 20 mg shortened the seizure duration by about 1 min, and delayed seizures by 12 min, whereas the

Table I. Flavonoids – subclasses division.

Flavonoids				
Flavonols	Flavanones	Flavones	Isoflavones	Chalcones
quercetin	naringenin	apigenin	daidzein	phloretin
routine				
kemferol	naringin	diosmetin	genistein	arbutin
mirecitin	hesperidin	luteolin		phloridzin
fistein	hesperedine	baicalin		chalconaringenin
morina				

Table II. Antiepileptic properties of selected flavonoids in animal models of epilepsy – a summary of scientific reports.

Flavonoids	Animal model	Results	Author
Naringenine	PILO-model (mice)	• reduction of seizures	Shakeel et al. (2017)
	KA-model (mice/rats)	• reduction of seizures, undisturbed cognitive functions	Park et al. (2016) Golechha et al. (2011).
	PTZ-model (rats)	• protection against seizures, reduction of cognitive impairment	Golechha et al. (2014)
Apigenin	picROTOXIN-induced seizures (mice)	• delayed the onset of seizures	Avallone et al. (2000)
	KA-model (mice)	• reduction of the number of seizures	Han et al. (2012)
Luteolin	PTZ-model (mice/rats)	• delayed the onset of myoclonic jerks, onset of clonic seizures and onset of hind limb extension • protection against mortality • neuroprotective effect	Tambe et al. (2015) Zhen et al. (2016)
	PILO-model (mice)	• delayed the onset of seizures and shortened the duration of clonic seizures • improvement of cognitive functions	Smillin et al. (2020)
Epigallocatechin	PTZ- kindling model (rats)	• reduced cognitive deficits and oxidative damage	Xie et al. (2012)
	cocaine-induced seizures (rats)	• reduction of seizures	Park et al. (2001)
	lithium-PILO model of TLE (rats)	• reduction of the frequency of spontaneous recurring seizures • neuroprotective effect	Qu et al. (2019)
Hesperidin	PTZ-model (mice/zebrafish)	• increased anticonvulsant effect alone and in combination with diazepam • neuroprotective effect	Kumar et al. (2014) Rosa-Falero et al. (2015)
Quercetin/routine	PTZ-kindling model (rats/mice)	• attenuation of seizures • memory improvement	Nassiri-Asl et al. (2013) Singh et al. (2017)
	PTZ model (rats)	• reduced generalized seizure	Sefil et al. (2014)
	MEST (mice)	• reduced seizures	Nieoczym et al. (2014)
	6 Hz model (mice)	• reduced seizures • no side effects in behavioral tests in combination with LEV and VPA	Nieoczym et al. (2014)
Baicalin	PILO-model (rats)	• significant delay in seizures • reduced neuronal death	Liu et al. (2012)
	KA-model (mice)	• neuroprotective effect	Liao et al. (2016)

higher dose (40 mg/kg) shortened seizure duration by 2 min and delayed seizures by 27 min. Moreover, naringenin treatment restored the antioxidant status and reduced lipid peroxidation in hippocampus of epileptic mice.

Park et al. (2016) using a kainic acid (KA) mouse model of temporal lobe epilepsy (TLE) showed that treatment with naringenin (100 mg) significantly delayed the occurrence of KA-induced seizures and impaired granular cell dispersion (GCD). Moreover, the effects of naringenin treatment were similar to those of rapamycin (RA) treatment. RA is not a universal antiepileptic drug and its use is limited to tuberous scler-

osis. In these studies, it was used as a positive control for changes in GCD mediated by mTORC1 inactivation. Nissel staining showed that naringenin significantly reduced changes in GCD, suggesting that it may act by inhibiting mTORC1 pathway activation. Many studies have indicated that hyperactivation of the mTORC1 pathway leads to epilepsy (Laplante et al., 2012; Sha et al., 2012; Lasarge et al., 2014; Shima et al., 2015). Immunofluorescence results and Western blot analysis showed that naringenin treatment attenuated the increase in TNF α and IL-1 β expression and significantly reduced their levels within the dentate gyrus (DG) of the hippocampus of KA-treated mice. These results




Fig. 2. Structure of naringenin.

suggest that naringenin may have anti-inflammatory effects in the KA-induced TLE model.

Interestingly, a flavanone glycoside naringin was tested in the KA induced epilepsy model in rats (Golechha et al., 2011). Results they obtained indicated, that initial treatment with naringin at doses 20, 40, 80 mg/kg significantly increased seizure delay compared to the KA control group, which suggests that naringin suppresses KA-induced seizures. Behavioral studies in the passive avoidance test showed cognitive impairment in the KA control group, while pretreatment with naringin reversed the KA-induced cognitive deficit (Golechha et al., 2011). Additionally, in the rat pentylentetrazol (PTZ) model naringin (20, 40 and 80 mg/kg) administered for 7 days significantly delayed the occurrence of myoclonic seizures depending on the dose (Golechha et al., 2014). Obtained results indicated that treatment with naringin (80 mg/kg) protected all rats against PTZ seizures, and reduced brain malondialdehyde and TNF- α levels and conserved glutathione, which highlighted the antioxidant, anti-inflammatory and anticonvulsant potential of naringin. Additional studies were conducted in order to check whether the anticonvulsant effect of naringin is GABA dependent. For this purpose, the GABA-A receptor was blocked by flumazenil, an antagonist of GABA-A. Obtained results showed, that flumazenil pretreatment decreased the prolongation of convulsant latency induced by naringin, suggesting a possible modulation of the benzodiazepine site of the GABA-A receptor by naringin to produce its anticonvulsant effect (Golechha et al., 2014).

Compans and coworkers (2018) investigated a potent anticonvulsant properties of two methylated flavanones: naringenin 4',7-dimethyl ether (NRG-DM) and naringenin 7-O-methyl ether (NRG-M) in zebrafish pentylentetrazole (PTZ) seizure model. Results they obtained showed, that both NRG-M NRG-DM decreased PTZ-induced seizure activity in larval zebrafish up to approximately 20-30% compared to PTZ controls. In

addition, NRG-DM was shown to be active in two standard mouse models of acute epilepsy, the PTZ test and in the 6 Hz psychomotor seizure model, suggesting that methylation facilitates the absorption of the compound, which translates into higher anticonvulsant activity. NRG-DM certainly deserves further research in the treatment of generalized seizures, and especially in the treatment of drug-resistant focal seizures (Compans et al., 2018).

Apigenin

Apigenin (4',5,7-trihydroxyflavone, Fig. 3) is one of the most common flavonoids in plants and belongs to the flavone subclass. It has been shown, that the main source of the apigenin are plants belonging to *Asteraceae*, such as *Artemisia* (Ornano et al., 2016), *Achillea* (Venditti et al., 2015; 2016), *Matricaria* (Sharifi-Rad et al., 2018) and *Tanacetum* (Venditti et al., 2018). Apigenin has received particular interest because of its anxiolytic, sedative, neuroprotective, anticonvulsant and antidepressant properties (Ali et al., 2017). Interestingly, apigenin can also ameliorate memory dysfunction linked to Alzheimer's disease (Zhao et al., 2013).

Studies by Avallone et al. (2000) determining a potent anticonvulsant efficacy of apigenin in picrotoxin-induced seizures in Sprague-Dawley male rats indicated, that apigenin at doses of 25 and 50 mg/kg administered 15 min before picrotoxin (6 and 8 mg/kg) delayed the onset of convulsions.

Interestingly, apigenin administrated for five days (25-50 mg/kg) followed by a KA injection on the last day, was shown to reduce both the number and duration of seizures in mice (Han et al., 2012). Moreover, apigenin was found to block KA-induced electroencephalogram discharge activity in the cerebral cortex. Cresol violet staining of the brain sections of KA-treated mice showed that KA causes the loss of approximately 20% of neurons, whereas apigenin administration af-




Fig. 3. Structure of apigenin.

ter KA treatment for 5 days rescued about 32-40% of the neuronal loss in the CA3 region. Additionally, apigenin (25-50 mg) in dose-dependent manner reversed the reduction of glutathione (GSH) levels induced by KA in hippocampus tissues. *In vitro* studies on primary hippocampal cell culture indicated that apigenin inhibited KA-induced excitotoxicity in a dose-dependent manner as measured by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) test. Moreover, apigenin (5, 20, 60 μ M) treatment dose-dependently inhibited intracellular reactive oxygen species (ROS) elevations in cultured hippocampal neurons. Other *in vitro* studies using SH-SY5Y cells indicated apigenin's neuroprotection against oxidative stress-induced cell death by blocking caspase-3 activity and glutamate-induced neurotoxicity by reducing NMDA receptor-mediated responses, which was demonstrated in cultured cortical neurons (Losi et al., 2004). In addition, apigenin was reported to inhibit the influx of extracellular Ca^{2+} and release of intracellular Ca^{2+} in the rat thoracic aorta and NMDA or gamma-aminobutyric acid (GABA) receptors channels (Ko et al., 1991).

Luteolin

Luteolin (3',4',5,7-tetrahydroxy flavone, Fig. 4) also belongs to the group of flavones. This compound is present in herbs such as thyme, chamomile, celery and green peppers. It can also be found in vegetables and fruits such as parsley, celery, broccoli, apples, onion leaves (Miean et al., 2001; Mencherini et al., 2007). Plants rich in luteolin have been used in the treatment of hypertension, inflammatory diseases and cancer (Harborne and Williams, 2000). The blood-brain barrier is permeable to luteolin, which makes it useful in the treatment of diseases of the central nervous system, including brain cancer (Wruck et al., 2007). Shaikh et al. (2013) indicated no anticonvulsant effect of doses (0.3-10 mg/kg) of luteolin in the 6 Hz seizure




Fig. 4. Structure of luteolin.

model, PTZ-induced seizure test, maximal electroshock (MES) tests as well as in PILO model with second hit PTZ seizure model in mice. On the other hand, Tambe and colleagues (2015) evaluated the antiepileptic potential of luteolin isolated from *Eclipta alba* in PTZ-induced seizure test in mice. Obtained results showed that the administration of luteolin at a dose of 10 or 20 mg/kg delayed the occurrence of myoclonic seizures, clonic seizures and extension of the hind limbs in comparison with the PTZ control animals. In addition, the above doses showed 100% protection against PTZ-induced mortality. The efficacy was comparable to that of diazepam, suggesting that luteolin is anti-epileptic in the PTZ-induced seizure test in a dose-dependent manner.

Zhen et al. (2016) studied the anticonvulsant effect of luteolin in the PTZ test in rats. They showed that initial treatment with luteolin at a dose of 100 mg/kg increased seizure delay, reduced the mean seizure duration and seizure severity compared to the PTZ group. Mortality in the PTZ group was 33.3%, while in the PTZ+100 mg/kg luteolin rats it decreased to 8.3%. Despite the fact that luteolin did not completely inhibit seizures, the results of Nissel's hippocampus staining and the Morris water maze behavioral test showed a significant reduction in the neurological damage and cognitive deficits associated with daily seizures. Biochemical studies indicated that the neuroprotective effect of luteolin on seizure-induced cognitive dysfunction was exerted by inhibition of ROS production and increased phosphoactivation of the cyclic AMP binding protein (CREB) pathway and expression of derived neurotrophic factor (BDNF) in the hippocampus (Zhen et al., 2016).

Interesting results were shown by Smilin Bell Aseervatham (2020) studying *Passiflora caerulea* (PCE), which contains polyphenol compounds such as ginsenoside, naringenin, chrysoeriol 8-c-glucoside, luteolin-6-C-glucoside, apigenin-6,8-di-C- β -D-glucopyranoside. PCE (200 mg/kg) significantly delayed the onset of seizures and reduced the duration of clonic seizures compared to the PCE group (100 mg/kg) and diazepam in the mouse PILO model of epilepsy. Furthermore, PCE-treated mice showed an improvement in cognitive function. It is likely that the antioxidant substances contained in PCE have a protective effect on PILO-induced seizures by neutralizing oxidative damage (Smilin Bell Aseervatham et al., 2020).

Epigallocatechin

Epigallocatechingallate (EGCG, 2R,3R-3',4',5,5',7-pentahydroxy flavan-3-yl gallate, Fig. 5) is classified in hydroxylated flavans. It constitutes 50% of all polyphenols

indicating an interaction with glutamate receptors, in particular with NMDA receptors and mGluR II, which suggests that the extract may be an antagonist at the glutamate binding site (Rosa-Falero et al., 2015).

Quercetin

Quercetin (3,3',4,4',7-pentahydroxy flavone, Fig. 7), a natural flavonoid classified as flavonols. Quercetin is found in the apples, blueberries, nuts, grapes, onions or chamomile (Srivastava et al., 2010). Nassiri-Asl et al. (2013) investigated the possible effects of quercetin on oxidative stress and memory retrieval in rat PTZ kindling model. Obtained results indicated that pre-treatment with 50 mg/kg of quercetin attenuated seizure severity from the beginning of the kindling procedure by lowering the mean seizure stages compared to the control group. Moreover, quercetin at dose of 50 mg/kg significantly increased the step-through latency of the passive avoidance response compared to the control rats.

Nieoczym et al. (2014) investigated the effects of routine and quercetin in some selected mouse epilepsy models, (6 Hz seizure, maximal electroshock seizure threshold MEST and intravenous (iv) PTZ threshold seizure tests). Rutine and quercetin (400 mg/kg) showed statistically significant anticonvulsant effect in the 6 Hz seizure test. There was no effect on thresholds in MEST and iv PTZ tests in mice. Moreover, they evaluated the effect of quercetin on the anticonvulsant effect of two AEDs, i.e., valproic acid (VPA) and levetiracetam (LEV) in a 6 Hz mouse model. Obtained results showed no significant impact of quercetin on the AEDs anticonvulsant efficacy. Similarly, no side effects in behavior (long-term memory, muscle strength or motor coordination) were observed.




Fig. 7. Structure of quercetin.

Sefil et al. (2014) showed that quercetin administration (10 and 20 mg/kg i.p.) reduced generalized seizure duration in PTZ (45 mg/kg i.p.) induced seizures in rats, whereas in picrotoxin induced seizures was ineffective. Quercetin also significantly decreased the seizure severity score and prolonged the onset of seizure at 10 mg/kg dose in PTZ induced seizure.

Furthermore, quercetin combined with LEV dose dependently attenuated depressive behavior in PTZ kindling mice (Singh et al., 2017). Long-term treatment with LEV without quercetin significantly reduced seizure severity, but exacerbated depressive behavior in mice with epilepsy and comorbid depression. This is due to the facts that LEV treatment of PTZ mice alters the kynureuin (KYN) and tryptophan (TRP) ratio in both cortical and hippocampal areas of the mouse brain, while LEV + quercetin treatment restores KYN:TRP levels to normal (Singh et al., 2017).

Baicalin

Baicalin ((2S,3S,4S,5R,6S)-6-(5,6-dihydroxy-4-oxo-2-phenyl-chromen-7-yl)oxy-3,4,5-trihydroxy-tetrahydropyran-2-carboxylic acid, Fig. 8), is a flavone glycoside found in the plant species *Scutellaria*. It has the following pharmacological properties: anti-tumor, antioxidant, antibacterial (Huang et al., 2019). Research by Liu et al. (2012), indicated anticonvulsant and neuroprotective effects of this compound on pilocarpine-induced epileptic model in rats. They showed that pretreatment baicalin (100 mg) delayed PILO-induced seizures in rats as well as decreased animal mortality. Additionally, baicalin reversed the changes in oxidative stress levels in the hippocampus (significantly reduced nitrite, glutathione and lipid peroxidation) compared to the PILO group. Immunohistochemical studies (Nis-




Fig. 8. Structure of baicalin.

sel, Fluoro-Jade B and TUNEL staining) showed that baicalin protected the hippocampus against neuronal loss, apoptosis and pilocarpine-induced degeneration. Liao et al. (2016) demonstrated anti-apoptotic properties of baicalin and proved that it protects against damage to the hippocampus in mice after KA-induced SE. Research indicates a decrease in the expression of miR-49 responsible for neuronal death and an increase in the level of Bcl-2 (an anti-apoptotic protein). It can be concluded that baicalin has a protective effect in SE and could become a potential supplement in brain damage prevention and, with further research, the treatment of epilepsy.

Toxicity studies

Since flavonoids are of great interest due to their broad spectrum of activity, they have also been tested for toxicity. Liang et al. (2017) assessed the toxicity of the flavonoid fraction from *Lithocarpus polystachyus* in rodents. In the case of acute toxicity, doses of 5000 mg/kg were used, and in the case of sub-chronic toxicity, doses were established at 0, 70, 140, 560 mg/kg for 26 weeks. The dose of 5000 mg/kg caused no mortality or any significant toxicity symptoms. In the sub-chronic toxicity experiment in rats no behavioral changes, convulsions, coma, or eating problems were observed. Analysis of blood coagulation parameters and biochemical parameters (creatinine, alkaline phosphatase, glucose, total protein, alanine aminotransferase, aspartate aminotransferase, bilirubin, total cholesterol, triglycerides) showed no significant changes in comparison to the control. Hematology only showed an increase in white blood cell (WBC) levels compared to the control. WBC levels were still normal. Increasing the level of WBC can help to strengthen the immune function of the body. Histopathological examinations did not reveal any pathological changes within organs. Peng and coworkers (2016) studied the toxicity of corn silk flavonoids in mice. General clinical signs, mortality, hematological, biochemical and histopathological parameters were examined. The analysis of the results did not show any significant changes compared to the control group. Research shows that the use of corn silk as functional food is safe.

CONCLUSIONS

Despite high availability of several generations of AEDs, patients resistant to a one drug treatment only require an additional anticonvulsant, which carries

the risk of increasing side effects. Flavonoids were shown to have a strong pharmacological effect and a wide spectrum of activity. In the light of the results of numerous *in vivo* studies, there is no doubt that due to the prevalence of flavonoids in the plant world and the presence of vegetables and fruits in the diet, these compounds are of great importance in the prevention of many civilization diseases and could prove effective as an add-on therapy for epilepsy. Thus, flavonoids can be useful for developing new therapeutic strategies for treating epilepsy. Although antiepileptic effects of flavonoids have been verified in preclinical studies, there is a great need for including flavonoids as antiepileptic agents in clinical trials.

REFERENCES

- Ali F, Rahul, Naz F, Jyoti S, Siddique YH (2017) Health functionality of apigenin: A review. *Int J Food Prop* 20: 1197–1238.
- Avallone R, Zanolli P, Puia G, Kleinschnitz M, Schreiber P, Baraldi M (2000) Pharmacological profile of apigenin, a flavonoid isolated from *Matricaria chamomilla*. *Biochem Pharmacol* 59: 1387–1394.
- Boon P, Ryvlin P, Wheless JW, Kawai K (2015) Treating drug-resistant epilepsy – why are we waiting? *Eur Neurol Rev* 10: 171–175.
- Bravo L (1998) Polyphenols: Chemistry, dietary, sources, metabolism, and nutritional significance. *Nutr Rev* 56: 317–333.
- Cano A, Ettcheto M, Chang JH, Barroso E, Espina M, Kühne BA, Barenys M, Auladell C, Folch J, Souto EB, Camins A, Turowski P, García ML (2019) Dual-drug loaded nanoparticles of Epigallocatechin-3-gallate (EGCG)/Ascorbic acid enhance therapeutic efficacy of EGCG in a APPswe/PS1dE9 Alzheimer's disease mice model. *J Control Release* 301: 62–75.
- Chen R, Qi QL, Wang MT, Li QY (2016) Therapeutic potential of naringin: an overview. *Pharm Biol* 54: 3203–3210.
- Cho J (2006) Antioxidant and neuroprotective effects of hesperidin and its aglycone hesperetin. *Arch Pharm Res* 29: 699–706.
- Cicero AFG, Fogacci F, Bove M, Giovannini M, Borghi C (2019) Three-arm, placebo-controlled, randomized clinical trial evaluating the metabolic effect of a combined nutraceutical containing a bergamot standardized flavonoid extract in dyslipidemic overweight subjects. *Phytother Res* 33: 2094–2101.
- Compans D, Orellana-Paucar AM, Steurs G, Zhang Y, Ny A, Foubert K, Exarchou V, Siekierska A, Kim Y, De Borggraave W, Dehaen W, Pieters L, de Witte PAM (2018) Methylated flavonoids as anti-seizure agents: Naringenin 4',7-dimethyl ether attenuates epileptic seizures in zebrafish and mouse models. *Neurochem Int* 112: 124–133.
- Corsale I, Carrieri P, Martellucci J et al., (2018) Flavonoid mixture (diosmin, troxerutin, rutin, hesperidin, quercetin) in the treatment of I-III degree hemorrhoidal disease: a double-blind multicenter prospective comparative study. *Int J Colorectal Dis* 33: 1595–1600.
- Del Rio D, Stewart AJ, Mullen W, Burns J, Lean ME, Brighenti F, Crozier A (2004) HPLC-MSn analysis of phenolic compounds and purine alkaloids in green and black tea. *J Agric Food Chem* 52: 2807–2815.
- Goldwasser J, Cohen PY, Lin W, Kitsberg D, Balaguer P, Polyak SJ, Chung RT, Yarmush ML, Nahmias Y (2011) Naringenin inhibits the assembly and long-term production of infectious hepatitis C virus particles through a PPAR-mediated mechanism. *J Hepatol* 55: 963–971.
- Golechha M, Chaudhry U, Bhatia J, Saluja D, Arya DS (2011) Naringin protects against kainic acid-induced status epilepticus in rats: evidence for an antioxidant, anti-inflammatory and neuroprotective intervention. *Biol Pharm Bull* 34: 360–365.

- Golechha M, Sarangal V, Bhatia J, Chaudhry U, Saluja D, Arya DS (2014) Naringin ameliorates pentylenetetrazol-induced seizures and associated oxidative stress, inflammation, and cognitive impairment in rats: possible mechanisms of neuroprotection. *Epilepsy Behav* 41: 98–102.
- Han JY, Ahn SY, Kim CS, Yoo SK, Kim SK, Kim HC, Hong JT, Oh KW (2012) Protection of apigenin against kainate-induced excitotoxicity by anti-oxidative effects. *Biol Pharm Bull* 35: 1440–1446.
- Harborne JB, Williams CA (2000) Advances in flavonoid research since 1992. *Phytochemistry* 55: 481–504.
- He J, Xu L, Yang L, Wang X (2018) Epigallocatechin gallate is the most effective catechin against antioxidant stress via hydrogen peroxide and radical scavenging activity. *Med Sci Monit* 24: 8198–8206.
- Jadeja RN, Devkar RV (2014) Polyphenols and flavonoids in controlling non-alcoholic steatohepatitis. In: *Polyphenols in Human Health and Disease*; Academic Press: San Diego, CA, USA, p. 615–623.
- Kalilani L, Sun X, Pelgrims B, Noack-Rink M, Villanueva V (2018) The epidemiology of drug-resistant epilepsy: A systematic review and meta-analysis. *Epilepsia* 59: 2179–2193.
- Kawai M, Hirano T, Higa S, Arimitsu J, Maruta M, Kuwahara Y, Ohkawara T, Hagihara K, Yamadori T, Shima Y, Ogata A, Kawase I, Tanaka T (2007) Flavonoids and related compounds as anti-allergic substances. *Allergol Int* 56: 113–123.
- Khalatbary AR, Khademi E (2018) The green tea polyphenolic catechin epigallocatechin gallate and neuroprotection. *Nutr Neurosci* 23: 281–294.
- Kirienko A, Radak D, Maggioli A (2019) Clinical efficacy of once-daily micronized purified flavonoid fraction 1000 mg tablet in patients with symptomatic chronic venous disease. *Curr Med Res Opin* 35: 553–557.
- Ko FN, Huang TF, Teng CM (1991) Vasodilatory action mechanisms of apigenin isolated from *Apium graveolens* in rat thoracic aorta. *Biochim Biophys Acta* 1115: 69–74.
- Kozera-Kępnia A, Jastrzębski K, Klimek A (2013) Pharmacogenetic determinants of drug resistance in epilepsy (in Polish). *Aktualn Neurol* 13: 96–102.
- Kumar A, Lalitha S, Mishra J (2014) Hesperidin potentiates the neuroprotective effects of diazepam and gabapentin against pentylenetetrazole-induced convulsions in mice: Possible behavioral, biochemical and mitochondrial alterations. *Indian J Pharmacol* 46: 309–315.
- Kwan P, Arzimanoglou A, Berg AT, Brodie MJ, Allen Hauser W, Mathern G et al., (2010) Definition of drug resistant epilepsy: consensus proposal by the ad hoc Task Force of the ILAE Commission on Therapeutic Strategies. *Epilepsia* 51: 1069–1077.
- Laplante M, Sabatini DM (2012) mTOR signaling in growth control and disease. *Cell* 149: 274–293.
- Lasarge CL, Danzer SC (2014) Mechanisms regulating neuronal excitability and seizure development following mTOR pathway hyperactivation. *Front Mol. Neurosci* 7: 18.
- Liang J, Chen SX, Huang S et al. (2017) Evaluation of toxicity studies of flavonoid fraction of *Lithocarpus polystachyus* Rehd in rodents. *Regul Toxicol Pharmacol* 88: 283–290.
- Liao Z, Liang R, Shi S, Wang C, Yang W (2016) Effect of baicalin on hippocampal damage in kainic acid-induced epileptic mice. *Exp Therapeutic Med* 12: 1405–1411.
- Lin YS, Tsai YJ, Tsay JS, Lin JK (2003) Factors affecting the levels of tea polyphenols and caffeine in tea leaves. *J Agric Food Chem* 51: 1864–1873.
- Liu YF, Gao F, Li XW, Jia RH, Meng XD, Zhao R, Jing YY, Wang Y, Jiang W (2012) The anticonvulsant and neuroprotective effects of baicalin on pilocarpine-induced epileptic model in rats. *Neurochem Res* 37: 1670–1680.
- Losi G, Puia G, Garzon G, de Vuono MC, Baraldi M (2004) Apigenin modulates GABAergic and glutamatergic transmission in cultured cortical neurons. *Eur J Pharmacol* 502: 41–46.
- Mencherini T, Picerno P, Scesa C, Aquino R (2007) Triterpene, antioxidant, and antimicrobial compounds from *Melissa officinalis*. *J Nat Prod* 70: 1889–1894.
- Miean KH, Mohamed S (2001) Flavonoid (myricetin, quercetin, kaempferol, luteolin, and apigenin) content of edible tropical plants. *J Agric Food Chem* 49: 3106–3112.
- Nassiri-Asl M, Moghbelinejad S, Abbasi E, Yonesi F, Haghghi MR, Lotfizadeh M, Bazahang P (2013) Effects of quercetin on oxidative stress and memory retrieval in kindled rats. *Epilepsy Behav* 28: 151–155.
- Nieoczym D, Socała K, Raszewski G, Właż P (2014) Effect of quercetin and rutin in some acute seizure models in mice. *Prog Neuropsychopharmacol Biol Psychiatry* 54: 50–58.
- Opeskin K, Berkovic SF (2003) Risk factors for sudden unexpected death in epilepsy: a controlled prospective study based on coroners cases. *Seizure* 12: 456–464.
- Ornano L, Venditti A, Donno Y, Sanna C, Ballero M, Bianco A (2016) Phytochemical analysis of non-volatile fraction of *Artemisia caerulescens* subsp. *Densiflora* (Viv.) (Asteraceae), an endemic species of La Maddalena Archipelago (Sardinia-Italy). *Nat Prod Res* 30: 920–925.
- Panche AN, Diwan AD, Chandra SR (2016) Flavonoids: an overview. *J Nutr Sci* 5: e47.
- Park J, Jeong KH, Shin WH, Bae YS, Jung UJ, Kim SR (2016) Naringenin ameliorates kainic acid-induced morphological alterations in the dentate gyrus in a mouse model of temporal lobe epilepsy. *Neuroreport* 27: 1182–1189.
- Park KK, Reuben JS, Soliman KFA (2001) The role of inducible-nitric oxide in cocaine induced kindling. *Exp Biol Med* 226: 185–190.
- Peng KZ, Zhang SY, Zhou HL (2016) Toxicological evaluation of the flavonoid-rich extract from *Maydis stigma*: Subchronic toxicity and genotoxicity studies in mice. *J Ethnopharmacol* 192: 161–169.
- Pietta PG (2000) Flavonoids as antioxidants. *J Nat Prod* 63: 1035–1042.
- Pinho-Ribeiro FA, Zarpelon AC, Fattori V, Manchope MF, Mizokami SS, Casagrande R (2016) Naringenin reduces inflammatory pain in mice. *Neuropharmacology* 105: 508–519.
- Qu Z, Jia L, Xie T, Zhen J, Si P, Cui Z, Xue Y, Sun C, Wang W (2019) Epigallocatechin-3-gallate protects against lithium-pilocarpine-induced epilepsy by inhibiting the toll-like receptor 4 (TLR4)/nuclear factor- κ B (NF- κ B) signaling pathway. *Med Sci Monit* 25: 1749–1758.
- Rayidi S, Pari L (2011) Effect of naringenin on carbohydrate metabolism in streptozotocin- nicotinamide induced diabetic rats. *Biomirror* 2: 12–19.
- Renugadevi J, Prabu SM (2009) Naringenin protects against cadmium-induced oxidative renal dysfunction in rats. *Toxicology* 256: 128–134.
- Rosa-Falero C, Torres-Rodríguez S, Jordán C, Licier R, Santiago Y, Toledo Z, Santiago M, Serrano K, Sosa J, Ortiz JG (2015) Citrus aurantium increases seizure latency to PTZ induced seizures in zebrafish thru NMDA and mGluR's I and II. *Front Pharmacol* 5: 284.
- Santos KF, Oliveria TT, Nagem TJ, Pinto AS, Oliveria MG (1999) Hypolipidemic effects of naringenin, rutin nicotinic acid and their association. *Pharmacol Res* 40: 493–496.
- Sefil F, Kahraman I, Dokuyucu R, Gokce H, Ozturk A, Tutuk O, Aydin M, Ozkan U, Pinar N (2014) Ameliorating effect of quercetin on acute pentylenetetrazole induced seizures in rats. *Int J Clin Exp Med* 7: 2471–2477.
- Sha LZ, Xing XL, Zhang D, Yao Y, Dou WC, Jin LR (2012) Mapping the spatio-temporal pattern of the mammalian target of rapamycin (mTOR) activation in temporal lobe epilepsy. *PLoS One* 7: e39152.
- Shaikh MF, Tan KN, Borges K (2013) Anticonvulsant screening of luteolin in four mouse seizure models. *Neurosci Lett* 550: 195–199.
- Shakeel S, Rehman MU, Tabassum N, Amin U, Mir MUR (2017) Effect of naringenin (a naturally occurring flavanone) against pilocarpine-induced status epilepticus and oxidative stress in mice. *Pharmacogn Mag* 13: S154–S160.
- Sharifi-Rad M, Nazaruk J, Polito L, Morais-Braga MFB, Rocha JE, Coutinho HDM, Salehi B, Tabanelli G, Montanari C, Del Mar Contreras M et al. (2018) *Matricaria* genus as a source of antimicrobial agents: From farm to pharmacy and food applications. *Microbiol Res* 215: 76–88.

- Shima A, Nitta N, Suzuki F, Laharie AM, Nozaki K, Depaulis A (2015) Activation of mTOR signaling pathway is secondary to neuronal excitability in a mouse model of mesio-temporal lobe epilepsy. *Eur J Neurosci* 41: 976–988.
- Shorvon S, Tomson T (2011) Sudden unexpected death in epilepsy. *Lancet* 378: 2028–2038.
- Singh T, Kaur T, Goel RK (2017) Adjuvant quercetin therapy for combined treatment of epilepsy and comorbid depression. *Neurochem Int* 104: 27–33.
- Smilin Bell Aseervatham G, Abbirami E, Sivasudha T, Ruckmani K (2020) Passifloracaerulea L. fruit extract and its metabolites ameliorate epileptic seizure, cognitive deficit and oxidative stress in pilocarpine-induced epileptic mice. *Metab Brain Dis* 35: 159–173.
- Srivastava JK, Shankar E, Gupta S (2010) Chamomile: A herbal medicine of the past with bright future. *Mol Med Rep* 3: 895–901.
- Sudha K, Rao AV, Rao A (2001) Oxidative stress and antioxidants in epilepsy. *Clin Chim Acta* 303: 19–24.
- Surges R, Thijs RD, Tan HL, Sander JW (2009) Sudden unexpected death in epilepsy: risk factors and potential pathomechanisms. *Nat Rev Neurol* 5: 492–504.
- Tambe R, Patil A, Jain P, Sancheti J, Somani G, Sathaye S (2017) Assessment of luteolin isolated from *Eclipta alba* leaves in animal models of epilepsy. *Pharm Biol* 55: 264–268.
- Tang F, Hartz AMS, Bauer B (2017) Drug-resistant epilepsy: multiple hypotheses, few answers. *Front Neurol* 8: 301.
- Tomson T, Nashef L, Ryvlin P (2008) Sudden unexpected death in epilepsy: current knowledge and future directions. *Lancet Neurol* 7: 1021–1031.
- Venditti A, Frezza C, Sciubba F, Serafini M, Bianco A, Cianfaglione K, Maggi F (2018) Volatile components, polar constituents and biological activity of tansy daisy (*Tanacetum macrophyllum* (Waldst. et Kit.) Schultz Bip.). *Ind Crop Prod* 118: 225–235.
- Venditti A, Guarcini L, Bianco A, Rosselli S, Bruno M, Senatore F (2016) Phytochemical analysis of *Achillea ligustica* all. from Lipari Island (*Aeolian islands*). *Nat Prod Res* 30: 912–919.
- Venditti A, Maggi F, Vittori S, Papa F, Serrilli AM, Di Cecco M, Bianco A (2015) Antioxidant and α -glucosidase inhibitory activities of *Achillea tenorii*. *Pharm Biol* 53: 1505–1510.
- Vuong QV, Golding JB, Nguyen M, Roach PD (2010) Extraction and isolation of catechins from tea. *J Sep Sci* 33: 3415–3428.
- Wang TY, Li Q, Bi KS (2018) Bioactive flavonoids in medicinal plants: Structure, activity and biological fate. *Asian J Pharm Sci* 13: 12–23.
- Wruck CJ, Claussen M, Fuhrmann G, Romer L, Schulz A, Pufe T, Waetzig V, Peipp M, Herdegen T, Gotz ME (2007) Luteolin protects rat PC12 and C6 cells against MPP+ induced toxicity via an ERK dependent Keap1-Nrf2-ARE pathway. *J. Neural Transm Suppl* 72: 57–67.
- Xie T, Wang WP, Mao ZF, Qu ZZ, Luan SQ, Jia LJ, Kan MC (2012) Effects of epigallocatechin-3-gallate on pentylentetrazole-induced kindling, cognitive impairment and oxidative stress in rats. *Neurosci Lett* 516: 237–241.
- Yokoi I, Kabuto H, Akiyama K, Mori A, Ozaki M (1989) Tannins inhibit the occurrence of epileptic focus induced by FeCl₃ injection in rats. *Jpn J Psychiatry Neurol* 43: 552–553.
- Zhao L, Wang JL, Liu R, Li XX, Li JF, Zhang L (2013) Neuroprotective, anti-amyloidogenic and neurotrophic effects of apigenin in an Alzheimer's disease mouse model. *Molecules* 18: 9949–9965.
- Zhen JL, Chang YN, Qu ZZ, Fu T, Liu JQ, Wang WP (2016) Luteolin rescues pentylentetrazole-induced cognitive impairment in epileptic rats by reducing oxidative stress and activating PKA/CREB/BDNF signaling. *Epilepsy Behav* 57: 177–184.
- Zhou W, Chen L, Hu X, Cao S, Yang (2019) Effects and mechanism of epigallocatechin-3-gallate on apoptosis and mTOR/AKT/GSK-3 β pathway in substantia nigra neurons in Parkinson rats. *Neuroreport* 30: 60–65.